
Assessing Stormwater Volume Reduction by Urban Trees















Urban Stormwater








Trees are an increasingly important part of stormwater management



"...inadequate research quantifying the urban tree contribution to rainfall/runoff processes limits their promotion by stormwater managers" Kuehler et. al., 2016

cubic feet of stormwater

California Central Valley – For every 1,000 trees,
stormwater is reduced by 1 million gallons



https://nepis.epa.gov/Exe/ZyPDF.cgi/P100H2RQ.PDF?Dockey=P100H2RQ.PD




Runoff Reduction by Trees: Highly Variable and Difficult to Measure



"The standard conversion factor of 59 gallons per tree per year and 18 gallons per tree seedling and shrub established will be used for purposes of review"


Final report will use i-Tree or
National Tree Benefit Calculator






Great Lakes Restoration Initiative - Lake Michigan Drainage Basin Fond du Lac, WI

This study will use a paired-basin approach to characterize the impact of tree removal on stormwater runoff characteristics from two medium-density residential catchments





Trees as Part of the Urban Water Balance



https://www.treesmatter.org/STBlog/3250 197

- Measure components of the urban water balance
- Apply measurements to improve models (i-Tree)












Cumulative Distribution of Runoff Volume







Predicted vs. Observed Runoff Volumes






 $1,636 \text{ m}^3 = 19\% \text{ of total runoff}$

2,990 m² of canopy = 574 liters/m²

31 trees removed = 52,774 liters/tree






432,185 gal = 19% of total runoff

32,184 ft² of canopy = 14 gal/ft^2


31 trees removed = 13,941 gal/tree





Comparison to Previous Studies

Location	Annual Per Tree (m³)	Interception (%)	Method
Fond du Lac, WI	52.7	43	Measured
Vancouver, BC		49-61	Measured
Melbourne, Aus.		29-44	Measured
Oakland, CA		14-27	Measured
Santa Monica, CA	6.6	27-65	Modeled
Modesto, CA	3.2		Modeled
Montgomery County, MD	7.6		Modeled
Cincinnati, OH	6.7		Modeled




- High variability among diversity of species, age, structure, etc.
- Annualized volume reduction per tree may be too general
 - Volume reduction as percent of annual precipitation is better
 - Volume reduction per unit area of canopy worth exploring



U.S. Forest Service: i-Tree Modeling

- Calibrate i-Tree Hydro using hyperlocal weather and discharge data
- Simulate discharge with and without tree cover
- Validate model predictions with post-tree removal discharge observations
- Assess performance and identify necessary improvements to i-Tree








Figure 2. GIS assessment of topography

Table 1. Preliminary estimate of change in surface runoff with loss of trees

of trees			
Basin Name	Preliminary Estimate of Change in Runoff if All Trees are Removed ^a		
Holly Tree lane Basin	+93.5 m ³ (+1.32% of base case surface runoff)		
Birch Tree Lane Basin	+186.7 m ³ (+1.36% of base case surface runoff)		

^a Difference in surface runoff if current tree canopy % in each area is converted to the land cover that was beneath that canopy, either herbaceous or impervious cover.



For more information: wrselbig@usgs.go

V





